REFEREANCES:

Augenbraun M, Rolfs R, Johnson R, Joesoef R, Pope V (1998). *Treponemal* specific tests for the serodiagnosis of syphilis. Sex Trans Dis. 25(10):549-52.

Bala K, Chaudhary U, Gill P, Mittal S, Vohra P (2013a). Seroprevalence of syphilis in HIV patients. Int J Pharm Med Bio Sc. 2(1):38-44.

Bala M, Singh V, Murlidhar S, Ramesh V (2013b). Assessment of reactivity of three treponemal tests in non-treponemal non-reactive cases from sexually transmitted diseases clinic, antenatal clinic, integrated counselling and testing centre, other different outdoor patient departments/indoor patients of a tertiary care centre and peripheral health clinic attendees. Indian J Med Microb. 31(3):275-79.

Binnicker M, Jespersen D, Rollins L (2012). Direct comparison of the traditional and reverse syphilis algorithm in a population with a low prevalence of syphilis. J ClinMicrobiol. 50(1):148-50.

Brown D, Frank J (2003). Diagnosis and management of syphilis. Am Fam Physician. 68(2):283-90.

Castro A, Esfandiari J, Kumar S, Ashton M, Kikkert S, Park M, et al (2010). Novel point-of-care test for simultaneous detection of non-treponemal and treponemal antibodies in patients with syphilis. J ClinMicrobiol. 48(12):4615-19.

Castro A, Jost H, Cox D, Fakile Y, Kikkert S, Tun Y, et al (2013). A comparison of the analytical level of agreement of nine treponemal assays for syphilis and possible implications for screening algorithms. BMJ Open. 3:e003347.

Castro A, Wang H (2011). Modified cardiolipin and uses therefore. US Patent no. 7,888,043 B2.

Chopdekar K, Patil S, Joshi A, Chowdhary A (2014). Serodiagnosis of syphilis in HIV sero-reactive patients. Indian J Bas Appl Med Res. 3(3):108-10.

Creegan L, Bauer M, Samuel M, Klausner J, Liska S, Bolan G (2007). An evaluation of the relative sensitivities of the venereal disease research laboratory test and the *Trepnemapallidum*

particle agglutination test among patients diagnosed with primary syphilis. Sex Transm Dis. 34(12):1016-18.

Das S, Kumar H (2014). Sero-prevalence of syphilis among voluntary blood donors: an institutional study. Int J Bas Appl Med Sci. 4(1):198-203.

Deshpande S (1996). Enzyme immunoassay: from concept to product development. In: Enzyme and signal amplification system. New York: Chapman and Hall. pp 155-91.

Egglestone S, Turner A (2000). Serological diagnosis of syphilis. Commun Dis Public Health. 3(3):158-62.

Goh B (2005). Syphilis in adults. Sex Transm Infect. 81(6):448-52.

Gu W, Yang Y, Wang Q, Pan B, Guo W, Wu L, et al (2013). Comparing the performance of traditional non-treponemal tests on syphilis and non-syphilis serum samples. Int J STD AIDS. 24(12):919-25.

Jafari Y, Peeling R, Shivkumar S, Claessens C, Lawrence J, Pai N (2013). Are *Treponemapallidum*specific rapid and point-of-care tests for syphilis accurate enough for screening in resource limited settings? Evidence from meta-analysis. PLoS One. 8(2):1-8.

Kingston M, French P, Goh B, Goold P, Higgins S, Sukthankar A (2008). UK national guideline on the management of syphilis 2008. Int J STD AIDS. 19(11):729-40.

Larsen S, Steiner B, Rudolph A (1995). Laboratory diagnosis and interpretation of tests for syphilis. Clin Microbiol Rev. 8(1):1-21.

Lee K, Park H, Roh E, Park K, Park M, Song E (2012). Characterization of sera with discordant result from reverse sequence screening for syphilis. Biomed Res Int. 2013:269347.

Lipinsky D, Schreiber L, Kopel V, Shainberg B (2012). Validation of reverse sequence screening for syphilis. J Clin Micbiol. 50(4):1501.

Loeffelholz M, Binnicker M (2012). It is time to use treponema-specific antibody screening tests for diagnosis of syphilis. J Clinc Microbiol. 50(1):2-6.

Marra C, Tantalo L, Maxwell C, Ho E, Sahi S, Jones T (2012). The rapid plasma regain test cannot replace the venereal disease research laboratory test for neurosyphilis diagnosis. Sex Transm Dis. 39(6):453-57.

Nayak S, Acharjya B (2012). VDRL test and its interpretation. Indian J Dermatol. 57(1):3-8.

Naidu N, Bharucha Z, Sonawane v, Ahmed I (2012). Comparative study of treponemal and non-treponemal test for screening of blood donated at a blood center. Asian J Transfus Sci. 6(1):32-5.

Pedersen N, Ørum O, Mouritsen S (1987). Enzyme-linked immunosorbent assay for detection of antibodies to the veneral disease research laboratory (VDRL) antigen in syphilis. J ClinMicrobiol. 25(9):1711-16.

Pope V, Fears M, Morrill W, Castro A, Kikkert S (2000). Comparison of the serodiaTreponemapallidum particle agglutination, captia syphilis-G, and spirotek regain II tests with standard test techniques for diagnosis of syphilis. J ClinMicrobio. 38(7):2543-45.

Ratnam S (2005). The laboratory diagnosis of syphilis. Can J Infect Dis Med Microbiol. 16(1):45-51.

Seňa A, White B, Sparling P (2010). Novel *Treponemapallidum* serologic tests: a paradigm shift in syphilis screening for the 21st century. Clin Infect Dis. 51(6):700-8.

Stone D, Moheng M, rolih S, Sinor L (1997). Capture-S, a non-treponemal solid-phase erythrocyte adherence assay for serological detection of syphilis. J Clin Microbiol. 35(1):217-22.

Tong M, Lin L, Liu L, Zhang H, Huang S, Chen Y, et al (2013). Analysis of 3 algorithms for syphilis serodiagnosis and implications for clinical management. Clin Infect Dis. 58(8):1116-24.

Tsang R, Radnos S, Morshed M (2011). Laboratory diagnosis of syphilis: a survey to examine the range of tests used in Canada. Can J Infect Dis Med Microbiol. 22(3):83-7.

Turbadkar D, Mathur M, Gaikwad S (2007). Prevalence of syphilis among HIV-seroreactive patients. Indian J Sex Transm Dis. 28(2):91-3.

White T, Fuller S (1989). Visuwell regain, a non-treponemal enzyme-linked immunosorbent assay for the serodiagnosis of syphilis. J Clin Microbiol. 27(10):2300-04.

Young H (2000). Guidelines for serological testing for syphilis. Sex Transm Inf. 76(5):403-5.

Zhang W, Yen-Lieberman B, Means c, Kreller R, Walezky J, Daly T (2012). The impact of analytical sensitivity on screening algorithm for syphilis. Clin Chem. 58(6):1065-6.